

/  
leptin levels. After administration of Escherichia coli lipopolysaccharide (LPS), leptin gene expression and leptin levels were increased. These findings suggest that leptin levels may be one mechanism by which anorexia is induced during acute inflammatory conditions.

=> display history 11-120

ENTER (BRIEF) OR FULL:brief

```
(FILE 'MEDLINE, BIOSIS' ENTERED AT 18:06:42 ON 07 JUN 2000)
L1      2173 S (INSULIN RECEPTOR SUBSTRATE-1) OR (IRS-1)
L2      44900 S (INSULIN-LIKE GROWTH FACTOR) OR (IGF)
L3      2136 S (GROWTH-FACTOR RECEPTOR-ASSOCIATED BINDING PROTEIN) OR GRB2
L4      39 S LEPTIN AND L1
L5      277 S LEPTIN AND L2
L6      1 S LEPTIN AND L3
L7      313 S L4 OR L5 OR L6
L8      205 DUP REM L7 (108 DUPLICATES REMOVED)
L9      107 S L8 AND PY<1999
L10     388 S (GAB)
L11     0 S LEPTIN AND L10
L12     97 S (GAB1 OR GAB2)
L13     1 S LEPTIN AND L12
L14     0 S L13 AND PY<1999
L15     215133 S (INTERLEUKIN OR (IL-4) OR (IL-9))
L16     169 S L15 AND LEPTIN
L17     113 DUP REM L16 (56 DUPLICATES REMOVED)
L18     32075 S (INTERLEUKIN-4) OR (INTERLEUKIN-9) OR (IL-4) OR (IL-9)
L19     5 S L18 AND LEPTIN
L20     3 DUP REM L19 (2 DUPLICATES REMOVED)
```

=> log y

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	147.73	147.88

STN INTERNATIONAL LOGOFF AT 18:57:16 ON 07 JUN 2000

Connection closed by remote host

Leptin      OB protein  
Obese